



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Genetic evidence of the association of DEAH-box helicase 37 defects with 46,XY gonadal dysgenesis spectrum

Citation for published version:

Evilen Da Silva, T, Gomes, NL, Lerário, AM, Keegan, CE, Nishi, MY, Carvalho, FM, Vilain, E, Barseghyanm, H, Martinez-aguayo, A, Forclaz, MV, Papazian, R, Pedroso De Paula, LC, Costa, EC, Carvalho, LR, Jorge, AA, Elias, F, Mitchell, R, Frade Costa, EM, Mendonca, BB & Domenice, S 2019, 'Genetic evidence of the association of DEAH-box helicase 37 defects with 46,XY gonadal dysgenesis spectrum', *The Journal of Clinical Endocrinology & Metabolism (JCEM)*. <https://doi.org/10.1210/jc.2019-00984>

Digital Object Identifier (DOI):

[10.1210/jc.2019-00984](https://doi.org/10.1210/jc.2019-00984)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

The Journal of Clinical Endocrinology & Metabolism (JCEM)

Publisher Rights Statement:

This is a pre-copyedited, author-produced version of an article accepted for publication in [insert journal title] following peer review. The version of record is available online at: [10.1210/jc.2019-00984](https://doi.org/10.1210/jc.2019-00984)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Genetic evidence of the association of DEAH-box helicase 37 defects as a novel
cause of 46,XY gonadal dysgenesis spectrum

Thatiana Evilen da Silva PhD^{1*}, Nathalia Lisboa Gomes MD^{1*}, Antonio Marcondes Lerário
MD,^{2,3} Catherine Elizabeth Keegan MD,^{4,5} Mirian Yumi Nishi PhD,¹ Filomena Marino
Carvalho MD,⁶ Eric Vilain MD,⁷ Hayk Barseghyanm PhD,⁷ Alejandro Martinez-Aguayo
MD,⁸ María Verónica Forclaz MD,⁹ Regina Papazian MD⁹, Leila Cristina Pedroso de Paula
MD¹⁰, Eduardo Corrêa Costa MD,¹⁰ Luciani Renata Carvalho MD,¹ Alexander A Jorge
MD¹, Felipe Elias MS¹, Rod Mitchell, MBChB, PhD¹¹, Elaine Maria Frade Costa MD,¹
Berenice Bilharinho Mendonca MD,^{1,2*} Sorahia Domenice MD,^{1*}

¹Unidade de Endocrinologia do Desenvolvimento, Laboratório de Hormônios e Genética
Molecular (LIM/42) da Disciplina de Endocrinologia e Metabologia do Hospital das Clínicas
da Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brasil

²Laboratório de Sequenciamento em Larga Escala (SELA), Faculdade de Medicina da
Universidade de São Paulo, São Paulo, Brasil

³Division of Metabolism, Endocrinology and Diabetes, Department of Internal Medicine,
University of Michigan, Ann Arbor, MI, USA

⁴Department of Pediatric Genetics, University of Michigan Medical School, Ann Arbor, MI,
USA

⁵Department of Human Genetics, University of Michigan Medical School, Ann Arbor, MI,
USA

⁶Departamento de Patologia, Faculdade de Medicina, Universidade de São Paulo, São Paulo,
SP, Brasil

⁷Center for Genetic Medicine Research, Children's National Health System, Children's Research Institute, Children's National Medical Center, Washington, DC, USA

⁸Division de Pediatría, Escuela de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile

⁹Servicio de Pediatría, Hospital Nacional Prof. Dr. A. Posadas, Buenos Aires, Argentina

¹⁰Hospital de Clínicas de Porto Alegre, Programa de Atendimento às Desordens do Desenvolvimento Sexual (PADS), Brasil, UFRGS

¹¹MRC Centre for Reproductive Health, Queens Medical Research Institute, Edinburgh, UK

*First authors da Silva, TE and Gomes, NL and last authors Mendonca, BB and Domenice, S contributed equally to this article.

All authors have nothing to disclose.

Short title: *DHX37* defects in 46,XY Gonadal Dysgenesis

Key words: *DHX37*; 46,XY gonadal dysgenesis; embryonic testicular regression syndrome

Corresponding author: Berenice B Mendonca

Hospital das Clínicas, Faculdade de Medicina da Universidade de São Paulo, Disciplina de Endocrinologia e Metabologia.

Av. Dr. Enéas de Carvalho Aguiar, 155, 2º andar, bloco 6

CEP: 05403-900, São Paulo, Brasil.

Email: beremen@usp; sorahiad@gmail.com

Phone number: 55 11 2661-7564; 55 11 2661-7512

48

49

50 **Acknowledgements**

51 This work was supported by grants from the Conselho Nacional de Desenvolvimento Científico
52 e Tecnológico (CNPq, Grant No. 305743/2011-2), the Fundação de Amparo à Pesquisa do
53 Estado de São Paulo (FAPESP, Grants No. 05/04726-0, 07/512156, 10/51102-0, 2013/02162-
54 8 and 2014/50137-5), and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior
55 (CAPES/ PNPd; institucional fellowship Grant). The authors are very grateful to Dr. Frederico
56 Moraes Ferreira for his technical assistance with the *in silico* prediction analysis, Dr. Beverly
57 M. Yashar and Dr. John Park for medical care of patient F6:II-1, and Dr Ivo Jorge Prado
58 Arnhold for suggestions and significant review of the manuscript.

59

ABSTRACT

Context: 46,XY gonadal dysgenesis (GD) is a heterogeneous group of disorders with a wide phenotypic spectrum, including embryonic testicular regression syndrome (ETRS). Most patients with GD remain without a molecular diagnosis. **Objective:** To report a novel gene for 46,XY GD etiology, especially for ETRS. **Design:** Screening of familial cases of 46,XY GD using whole exome sequencing and sporadic cases by target gene panel sequencing. **Setting:** Tertiary referral center for Differences/Disorders of sex Development (DSD). **Patients and methods:** We selected 87 patients with 46,XY DSD (17 familial cases from eight unrelated families and 70 sporadic cases); 55 patients had GD (among them, ten patients from five families and eight sporadic cases had ETRS) and 32 patients had 46,XY DSD of unknown etiology. **Results:** We identified four heterozygous missense rare variants classified as pathogenic or likely pathogenic in *DEAH-box helicase 37 (DHX37)* gene in five families (n=11 patients) and in six sporadic cases. Two variants were recurrent: the p.Arg308Gln (in two families and in three sporadic cases) and the p.Arg674Trp (in two families and in two sporadic cases).

The variants were specifically associated with ETRS (7/14 index cases; 50%). The frequency of rare, predicted to be deleterious DHX37 variants in this cohort (0.14) is significantly higher than that observed in gnomAD population database (0.004; p<0.001). Immunohistochemistry analysis in human testis showed that DHX37 is mainly expressed in germ cells, at different stages of testis maturation, in Leydig cells and rarely in Sertoli cells. Conclusion: This strong genetic evidence identifies DHX37 as a new player in the complex cascade of male gonadal differentiation and maintenance.

Introduction

46,XY gonadal dysgenesis (GD) represents a heterogeneous group of disorders/differences of sex development (DSD) characterized by abnormal gonadal development leading to a wide phenotypic spectrum. Variable degrees of external genitalia undervirilisation are observed, ranging from micropenis to female-like genitalia and partially- or fully-developed Mullerian derivatives. The gonads from these patients display a wide spectrum of histological abnormalities, ranging from ovarian-like stroma with disorganized seminiferous tubules to complete absence of gonadal tissue (1). Embryonic testicular regression syndrome (ETRS) is considered a part of the clinical spectrum of 46,XY gonadal dysgenesis (2). Most individuals with ETRS present with micropenis or atypical genitalia and lack of gonadal tissue on one or both sides (2). Partial or complete Mullerian duct regression associated with micropenis suggests an intrinsically functional testis in the first months of fetal life subsequent loss of testicular function before the last trimester of gestation, when the increase in penile length occurs. Numerous genes are known to be involved in the process of gonadal determination (3). However, a genetic diagnosis is identified in less than 40% of the patients with 46,XY GD (4). Moreover, few patients with ETRS were included in large cohorts of 46,XY DSD previously studied (4). However, the fact that some familial cases of ETRS were reported indicates a genetic etiology (5,6). In the present work, high throughput parallel sequencing methods, including whole-exome sequencing (WES) and targeted DSD-gene panels, were used to investigate the underlying genetic etiology in a large cohort of 46,XY patients with GD and 46,XY DSD patients with unknown etiological cause.

We identified recurrent rare variants in DEAH (Asp-Glu-Ala-His) box polypeptide 37 (*DHX37*) in several affected individuals from distinct families, establishing a novel genetic cause for 46,XY gonadal dysgenesis spectrum, including ETRS.

Ethics

This study was approved by the Ethics Committee of the Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, the Institutional Review Board of the University of Michigan Medical School, the Hospital de Garrahan Escuela de Medicina, Pontificia Universidad Católica de Chile, and the Hospital Nacional Prof. Dr. A. Posadas, Buenos Aires, Argentina. Written informed consent was obtained from all patients, their parents or legal guardians.

Subjects and Methods

We studied eighty-seven 46,XY DSD patients without previous molecular diagnosis, including 17 familial cases of 46,XY GD from 8 non-consanguineous families and 70 sporadic cases (38 with GD and 32 with 46,XY DSD of unknown etiology). Out of the 55 patients with GD, 10 patients from 5 families and 8 sporadic cases had an ETRS phenotype. The patients had different nationalities: Brazilian (81 patients), Argentinian (three siblings), Chilean (two siblings) and Chinese-American (one patient).

All patients have a normal GTG-banded metaphases 46,XY karyotype.

The 46,XY DSD patients were classified as having complete GD (CGD) if they had female external genitalia, Mullerian derivatives and streak gonads; as partial GD (PGD) if they had atypical external genitalia, Mullerian derivatives and at least one gonad with histopathological features of dysgenetic testis; as ETRS if they had micropenis, partially developed Mullerian derivatives and no gonadal tissue or small area of gonadal stroma; and as 46,XY DSD of unknown etiology if hormonal profile was not conclusive or not available due to previous gonadectomy. In this latter group, molecular defects of LHCG

and androgen receptors, *CYP17A1*, *HSD17B3*, *HSD3B2*, and *5ARD2* genes were ruled out by DNA sequencing.

Genomic DNA

For molecular diagnosis, genomic DNA was extracted from peripheral blood leukocytes by the proteinase K-SDS salting-out method (7).

Genetic study

Whole exome sequencing (WES) was performed in 14 familial cases from 7 families. In all but one family, the probands and their first-degree relatives and other affected family members were studied.

Sixty-eight sporadic cases were studied by targeted massively parallel sequencing.

DHX37 was studied by Sanger sequencing in two sporadic cases and in three patients from Family 2 (Figure 1). (Supplementary information, in DOI: 10.13140/RG.2.2.35903.76968).

Enrichment for massively parallel sequencing was performed with Nextera Exome Enrichment Kit (Illumina, San Diego, CA), followed by paired-end sequencing on the Illumina HiSeq 2500 System (Illumina, San Diego, CA)

For target sequencing, we designed an amplicon-based capture panel against exonic regions of 63 genes, including 43 genes known to be associated with human DSDs and 20 candidate genes, including *DHX37* (3) (see Table S1, Supplementary information in DOI: 10.13140/RG.2.2.35903.76968). Target sequences were captured using a custom Sure Select Target Enrichment System Kit (Agilent Technologies, Santa Clara, CA, USA) and sequencing was performed on the Illumina MiSeq platform (San Diego, CA, USA).

Sanger sequencing was used to confirm the potentially pathogenic variants identified by massively parallel sequencing and for segregation analysis. Sequencing was performed

on the ABI 3730XL DNA Analyzer (Applied Biosystems) using the BigDye (Applied Biosystems), followed by data analysis using a Genetic Analyzer (ThermoFisher Scientific).

The identified variants were classified according to American College of Medical Genetics (ACMG) criteria (8).

Data analysis

The exome and the targeted panel sequencing data were screened for rare variants (minor allele frequency < 0.1% in the public databases: Genome Aggregation Database (gnomAD) (9), 1000 Genomes (10), and in the Brazilian population database (ABraOM) (11), located in exonic and consensus splice site regions. Subsequently, the filtration pipeline prioritized potentially pathogenic candidate variants (loss of function variants and variants classified as pathogenic by multiple *in silico* programs). For variants identified by WES, we selected variants that fitted an autosomal-dominant model.

The sequencing reads carrying candidate variants were visually confirmed using the Integrative Genomics Viewer (Broad Institute, Cambridge, MA). Candidate variants were segregated in the family members by Sanger method.

The filtering of the variants is provided in Supplementary data (Figure S1) DOI: 10.13140/RG.2.2.35903.76968)

Histological analysis

Immunohistochemical staining

Eight formalin-fixed paraffin-embedded testicular autopsy samples from 46,XY individuals with different chronological ages (27 and 33 weeks gestational age, 1, 53, and 180 days of age, 13, 23 and 53 years of age) were collected and used for DHX37 expression analysis by immunohistochemistry. All samples were sliced into 3-µm-thick sections using an automatic Leica RM2255 microtome (Leica Biosystems, Nussloch,

Germany). The sections were briefly stretched in xylol at 600°C for 20 min, cooled in xylol, and dried in an incubator (Fanem Orion 515, São Paulo, Brazil) at 600°C. Sections were subjected to hematoxylin-eosin (HE) staining for histological analysis. For immunohistochemical study, slides were deparaffinized with xylene, hydrated in ethanol, washed in phosphate-buffered saline (0.01 M/pH 7.4), and blocked using methanol and hydrogen peroxide. Epitope exposure was carried out by placing the slide in boiling 10 mM citric acid (pH 6) or 100 mM EDTA (pH 9), followed by blocking non-specific protein. Rabbit polyclonal anti-DHX37 antibody (NB110-40581; Novus Biologicals, USA) was added at a dilution factor of 1:50. Dilution was standardized after testing on ovarian and skin tissues where protein expression was identified in cytoplasm of oocytes and nuclear membranes of ovarian stromal and squamous cells. The samples were incubated with universal secondary antibodies using the Novo Link Detection Systems kit (Leica Biosystems, USA) according to the manufacturer's instruction.

Statistical analysis

To test the genetic evidence for the association between *DHX37* and GD phenotype, we performed aggregate variant analyses comparing allele frequencies among our 46,XY DSD cohort and public databases [gnomAD and ABraOM].

Variants with similar characteristics of the *DHX37* variants observed in our cohort (rare nonsynonymous variants with a minor allele frequency of 0.01 and located in the two highly conserved protein (ATP-binding and Helicase C-terminal domains) that are predicted to be pathogenic by at least four *in silico* tools (Mutation Taster, SIFT, PolyPhen-2, Mutation Assessor and PROVEAN) were selected. Allele frequency differences between groups were analyzed by X^2 test, and statistical significance was set at $p < 0.05$. Statistical analyses were performed using SIGMAstat statistical software package (Windows version 3.5; SPSS Inc., San Rafael, CA).

Results

Patient phenotype and *DHX37* variants

Firstly, WES identified the same *DHX37* variant p.Arg308Gln (c.923G>A) (GenBank: NM_032656.3) in heterozygous state in two unrelated Brazilian families with ETRS (Families 1 and 2). All the affected individuals have the same phenotype (micropenis and absence or bilateral rudimentary gonadal tissue) (Figure 1, Table 1). A founder effect for p.Arg308Gln variant was ruled out in Families 1 and 2.

The p.Arg308Gln variant was also identified by WES in a Chinese-American sporadic case of ETRS from Michigan University performed in Eric Vilain's laboratory (sporadic case F6:II-1, Figure 1, Table 2).

As a novel candidate gene for 46,XY DSD, *DHX37* was included in our target DSD-panel. The same p.Arg308Gln variant was identified in another two sporadic cases: one had ETRS (sporadic case F7:II-1) and the other had PGD (sporadic case F8:II-1) (Figure 1; Table 2).

A further three different heterozygous *DHX37* missense variants (the p.Arg674Trp, p.Ser595Phe and p.Thr304Met) were identified in seven affected members from three families and in three sporadic cases (Figure 2).

All of these four variants are predicted to be pathogenic by at least four *in-silico* prediction tools (Table 3) and are absent in genomic population databases, except for the p.Arg308Gln, which has a very low allele frequency (0.00003) in the gnomAD database (Tables 4-5).

The p.Arg674Trp (c.2020C>T) variant was identified in the two Chilean brothers, both with ETRS (cases F3:II-1 and F3:II-2, Family 3), and also in the three Argentinian affected members (two brothers with ETRS and their uncle with PGD; cases F4:III-1, F4:III-2 and F4:II-4, respectively, Family 4) (Figure 1, Table 1). In addition, the

p.Arg674Trp variant was also identified in another two Brazilian sporadic cases, one patient with ETRS (sporadic case F10:II-1) and the other with PGD (sporadic case F11:II-1) (Figure 1, Table 2)

The p.Ser595Phe (c.1784C>T) variant was identified in two affected individuals from the same Brazilian family (Family 5). The proband had PGD and her nephew had ETRS (F5:II-6 and F5:III-1, respectively) (Figure 1, Table 1).

The p.Thr304Met (c.911C>T) was identified in a Brazilian female (sporadic case F9: II-5), who had previously undergone bilateral gonadectomy and genitoplasty (Figure 1, Table 2).

The p.Arg308Gln variant is classified as pathogenic and the other three variants, p.Arg674Trp, p.Ser595Phe and p.Thr304Met, are classified as likely pathogenic accordingly the ACMG criteria (Tables 4-5).

Segregation analysis of *DHX37* variants

Segregation analysis of the *DHX37* variants in eight families displayed a sex-limited autosomal dominant pattern, maternally inherited in five families (F2, F3, F4, F5, F11). In the Family 1, the presence of the p.Arg308Gln variant in the asymptomatic father suggests an autosomal dominant pattern of inheritance with incomplete penetrance (Figure 1). In two sporadic cases (F6. II-1 and F8.II-1), the confirmed paternity displayed a *de novo* status of the p.Arg308Gln *DHX37* variant.

***DHX37* gene and its protein structure**

DHX37 is located in the 12q24.31 region. It is a member of the large DEAH family of proteins and encodes an RNA helicase (12). The *DHX37* protein (NP_116045) comprises 1157 amino acids and four main domains. The conserved motifs of the helicase core region contain the Helicase ATP-binding domain (position 262-429) and the Helicase superfamily c-terminal domain (position 585-674); the two other domains are the helicase

associated domain (position 768-859) and the oligonucleotide/oligosaccharide-binding-fold domain (position 894-1011) (Figure 2). All the identified variants are located in the helicase core region (Figure 2).

DHX37 protein was identified in different testicular cells

DHX37 expression was characterized in testes from newborns, children and adults using immunohistochemistry. DHX37 was expressed in fibroblasts, endothelial cells and epithelial cells of epididymis. These cells were used as internal positive controls for immunohistochemistry. We found DHX37 expression in Leydig cell cytoplasm and in germ cells at different stages of maturation. Our analysis indicates that DHX37 expression in spermatogonia is characterized by a regular perinuclear halo pattern in both newborns (five samples) and adults (three samples). This pattern of staining differs from that seen in Leydig cells (granular cytoplasmatic) and during other stages of maturation of germ cells. A progressive condensation of protein around the nucleus was observed as cells differentiate from spermatocytes to spermatids, generating a localized paranuclear dot-like pattern. There was no staining in spermatozoa. Rare Sertoli cells displayed a weak and focal cytoplasmatic stain (Figure 3).

Frequency of the *DHX37* variants in our 46,XY DSD cohort

The allele frequency of rare and predicted to be deleterious *DHX37* variants identified in our cohort of 46,XY DSD patients [11/78 index cases (0.14)] was markedly higher than that observed in individuals from gnomAD [568 /141456 individuals (0.004; $p<0.001$)] and from a Brazilian cohort [1/609 individuals (0.002); $p<0.001$].

Discussion

The present study analyzed a large cohort of 46,XY DSD patients without a molecular diagnosis, most of whom had a GD phenotype, including a large number of familial and

285 sporadic cases with ETRS.

286 Pathogenic or likely pathogenic allelic variants in the *DHX37* were identified in 11
287 familial cases from 5 unrelated families and in six sporadic cases. Deleterious variants
288 are recurrent in familial and sporadic cases of 46,XY GD in patients of different
289 nationalities.

290 The *DHX37* gene has never been directly associated with gonadal development, but
291 deletions or rearrangements of the 12q24 chromosomal region, which contains *DHX37*
292 gene, have been associated with atypical genital development (13). Four syndromic
293 patients with micropenis or perineal hypospadias, and/or hypergonadotropic
294 hypogonadism are reported to have deletions or rearrangements involving the 12q24
295 region (13-15).

296 The *DHX37* gene encodes a RNA helicase protein which is involved in RNA-related
297 processes, including transcription, splicing, ribosome biogenesis (16), translation and
298 degradation (12,17). *DHX37* is required for maturation of the small ribosomal subunit in
299 human cells, through its catalytic activity, required for dissociation of the U3 snoRNA
300 from pre-ribosomal complexes (18). Disturbance of human ribosome production is
301 associated with cancer and genetic diseases known as ribosomopathies (19).

302 Disease-causing variants in the DExH-box helicase 30 (*DHX30*), were previously
303 described in syndromic patients with global developmental delay, intellectual disability,
304 severe speech impairment and gait abnormalities. Functional studies of allelic variants in
305 *DHX30* demonstrated that they affect protein folding or stability interfering with the RNA
306 binding (mutations located in Motif Ia) or with ATPase activity (mutations located in
307 Motif II and VI) (17,18). Two *DHX37* allelic variants found in the present study are
308 located in the same motifs.

309 Despite lack of experimental evidence to formally demonstrate the deleterious effects of

the four variants identified in the present study, they are located in the highly conserved helicase core region of the DHX37 protein.

The spontaneous p.Leu489Pro Dhx37 pathogenic variant was identified in Zebrafish in association with behavior scape defects (20). This study demonstrated that Dhx37 is involved in pre-mRNA splicing reinforcing the role of Dhx37 in RNA-related processes. Although there is no direct evidence of *DHX37* being involved in mRNA processing during gonadal development, DExD/H-box RNA helicase genes are differentially expressed between males and females during the critical period of male sex differentiation in channel catfish (21).

Further, we show population evidence that the *DHX37* variants are enriched among the 46,XY GD patients in comparison with the population database. The statistical analysis confirmed that the predicted deleterious *DHX37* variants located in the helicase core region are more frequently identified in our 46,XY DSD cohort than in the public databases, emphasizing that this finding was not by chance ($p < 0.01$).

Therefore, *in vitro* and *in vivo* studies on DHX37 mechanism have demonstrated a role of DHX37 in ribosome biogenesis (23). Based on this new knowledge, 46,XY gonadal dysgenesis could be classified as a ribosomopathy, expanding the etiological mechanisms of dysgenetic 46,XY DSD spectrum.

Since the discovery of the sex-determining region Y (*SRY*) variants in patients with GD in 1990 (22), several genes have been associated with the molecular etiology of this disorder. The nuclear receptor subfamily 5 group A member 1 (*NR5A1*) and Mitogen-Activated Kinase Kinase Kinase 1 (*MAP3K1*) variants are the most frequent causes of 46,XY gonadal dysgenesis identified to date (23-26).

In this study we found pathogenic/ likely pathogenic variants in *DHX37* in patients with 46,XY GD at a frequency of 14%, which is slightly higher than the frequency of *NR5A1*

defects (11%) in our whole cohort (24,27). Considering only the ETRS phenotype (micropenis and absence of uni or bilateral testicular tissue) this frequency increases to 50% (7/14 families).

In the literature, different inheritance patterns have already been described in 46,XY gonadal dysgenesis kindreds (28), including the description of asymptomatic male carriers of proven pathogenic variants of genes involved in testicular determination, such as *SRY* and *NR5A1* genes (29,30). Uncertain mechanisms might prevent the appearance of the phenotype in asymptomatic 46,XY carriers.

Maternal inheritance was observed in all familial cases with pathogenic/ likely pathogenic variants in *DHX37* with the exception of family 1, where the variant was inherited from a seemingly unaffected father carrier.

In adult humans, the *DHX37* protein is expressed in the ovarian stroma and in the cells within seminiferous tubules (Human Protein Atlas database) (31-33). In our study, the immunohistochemistry analysis of normal testicular tissue from newborn, pubertal and adult males revealed that *DHX37* is expressed during specific stages of germ cell maturation, in Leydig cells and rarely in Sertoli cells.

An elaborate paracrine cell-cell network transporting signaling molecules between germ cells and Sertoli cells has been described (34). Indeed, *in vitro* studies have shown that there is a bidirectional trafficking between Sertoli and germ cells, and that each cell type regulates the function of the other (35-38). In addition, RNA expression profiles of *DHX37* in human testicular cancer cells are higher than in other tissues (The Human Protein Atlas – Pathology), suggesting that *DHX37* may be involved in the regulatory process of the cell proliferation in the testis (31-33).

The present study provides several lines of genetic evidence to indicate that defects in *DHX37* are associated with 46,XY GD spectrum, mainly with ETRS. First, we observed

that the variants segregate with the DSD phenotype in a dominant inheritance pattern in most of the families and that two *de novo* variants were identified. Second, we provide statistical evidence that rare *DHX37* variants are enriched in the analyzed 46,XY DSD cohort in comparison with public databases involving a large number of individuals not selected by this phenotype.

In conclusion, our findings indicate that *DHX37* is a new player in the complex cascade of male gonadal differentiation and maintenance, thus establishing a novel and frequent molecular etiology for 46,XY gonadal dysgenesis spectrum, which includes a high proportion of individuals with embryonic testicular regression syndrome.

Supplementary information: displayed in DOI: 10.13140/RG.2.2.35903.76968).

370 **References**

- 371 1. Robboy SJ, Miller T, Donahoe PK, Jahre C, Welch WR, Haseltine FP, Miller WA,
372 Atkins L, Crawford JD. Dysgenesis of testicular and streak gonads in the
373 syndrome of mixed gonadal dysgenesis: perspective derived from a
374 clinicopathologic analysis of twenty-one cases. *Hum Pathol.* 1982;13(8):700-716.
- 375 2. Marcantonio SM, Fechner PY, Migeon CJ, Perlman EJ, Berkovitz GD.
376 Embryonic testicular regression sequence: a part of the clinical spectrum of
377 46,XY gonadal dysgenesis. *Am J Med Genet.* 1994;49(1):1-5.
- 378 3. Ono M, Harley VR. Disorders of sex development: new genes, new concepts. *Nat*
379 *Rev Endocrinol.* 2013;9(2):79-91.
- 380 4. Eggers S, Sadedin S, van den Bergen JA, Robevska G, Ohnesorg T, Hewitt J,
381 Lambeth L, Bouty A, Knarston IM, Tan TY, Cameron F, Werther G, Hutson J,
382 O'Connell M, Grover SR, Heloury Y, Zacharin M, Bergman P, Kimber C, Brown
383 J, Webb N, Hunter MF, Srinivasan S, Titmuss A, Verge CF, Mowat D, Smith G,
384 Smith J, Ewans L, Shalhoub C, Crock P, Cowell C, Leong GM, Ono M, Lafferty
385 AR, Huynh T, Visser U, Choong CS, McKenzie F, Pachter N, Thompson EM,
386 Couper J, Baxendale A, Gecz J, Wheeler BJ, Jefferies C, MacKenzie K, Hofman
387 P, Carter P, King RI, Krausz C, van Ravenswaaij-Arts CM, Looijenga L, Drop S,
388 Riedl S, Cools M, Dawson A, Juniarto AZ, Khadilkar V, Khadilkar A, Bhatia V,
389 Dũng VC, Atta I, Raza J, Thi Diem Chi N, Hao TK, Harley V, Koopman P, Warne
390 G, Faradz S, Oshlack A, Ayers KL, Sinclair AH. Disorders of sex development:
391 insights from targeted gene sequencing of a large international patient cohort.
392 *Genome Biol.* 2016;17(1):243.
- 393 5. de Grouchy J, Gompel A, Salomon-Bernard Y, Kuttenn F, Yaneva H, Paniel JB,
394 Le Merrer M, Roubin M, Doussau de Bazignan M, Turleau C. Embryonic
395 testicular regression syndrome and severe mental retardation in sibs. *Ann Genet.*
396 1985;28(3):154-160.
- 397 6. Josso N, Briard ML. Embryonic testicular regression syndrome: variable
398 phenotypic expression in siblings. *J Pediatr.* 1980;97(2):200-204.
- 399 7. Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting
400 DNA from human nucleated cells. *Nucleic Acids Res.* 1988;16(3):1215.
- 401 8. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde
402 M, Lyon E, Spector E, Voelkerding K, Rehm HL, Committee ALQA. Standards
403 and guidelines for the interpretation of sequence variants: a joint consensus
404 recommendation of the American College of Medical Genetics and Genomics and
405 the Association for Molecular Pathology. *Genet Med.* 2015;17(5):405-424.
- 406 9. Brownstein CA, Beggs AH, Homer N, Merriman B, Yu TW, Flannery KC,
407 Dechene ET, Towne MC, Savage SK, Price EN, Holm IA, Luquette LJ, Lyon E,
408 Majzoub J, Neupert P, McCallie D, Jr., Szolovits P, Willard HF, Mendelsohn NJ,
409 Temme R, Finkel RS, Yum SW, Medne L, Sunyaev SR, Adzhubey I, Cassa CA,
410 de Bakker PI, Duzkale H, Dworzy Ski P, Fairbrother W, Francioli L, Funke BH,
411 Giovanni MA, Handsaker RE, Lage K, Lebo MS, Lek M, Leshchiner I, Macarthur
412 DG, McLaughlin HM, Murray MF, Pers TH, Polak PP, Raychaudhuri S, Rehm
413 HL, Soemedi R, Stitzel NO, Vestecka S, Supper J, Gugenmus C, Klocke B, Hahn
414 A, Schubach M, Menzel M, Biskup S, Freisinger P, Deng M, Braun M, Perner S,
415 Smith RJ, Andorf JL, Huang J, Ryckman K, Sheffield VC, Stone EM, Bair T,
416 Black-Ziegelbein EA, Braun TA, Darbro B, Deluca AP, Kolbe DL, Scheetz TE,
417 Shearer AE, Sompallae R, Wang K, Bassuk AG, Edens E, Mathews K, Moore
418 SA, Shchelochkov OA, Trapane P, Bossler A, Campbell CA, Heusel JW, Kwikite

A, Maga T, Panzer K, Wassink T, Van Daele D, Azaiez H, Booth K, Meyer N, Segal MM, Williams MS, Tromp G, White P, Corsmeier D, Fitzgerald-Butt S, Herman G, Lamb-Thrush D, McBride KL, Newsom D, Pierson CR, Rakowsky AT, Maver A, Lovre IL, Palanda IA, Peterlin B, Torkamani A, Wedell A, Huss M, Alexeyenko A, Lindvall JM, Magnusson M, Nilsson D, Stranneheim H, Taylan F, Gilissen C, Hoischen A, van Bon B, Yntema H, Nelen M, Zhang W, Sager J, Zhang L, Blair K, Kural D, Cariaso M, Lennon GG, Javed A, Agrawal S, Ng PC, Sandhu KS, Krishna S, Veeramachaneni V, Isakov O, Halperin E, Friedman E, Shomron N, Glusman G, Roach JC, Caballero J, Cox HC, Mauldin D, Ament SA, Rowen L, Richards DR, Lucas FA, Gonzalez-Garay ML, Caskey CT, Bai Y, Huang Y, Fang F, Zhang Y, Wang Z, Barrera J, Garcia-Lobo JM, Gonzalez-Lamuno D, Llorca J, Rodriguez MC, Varela I, Reese MG, De La Vega FM, Kiruluta E, Cargill M, Hart RK, Sorenson JM, Lyon GJ, Stevenson DA, Bray BE, Moore BM, Eilbeck K, Yandell M, Zhao H, Hou L, Chen X, Yan X, Chen M, Li C, Yang C, Gunel M, Li P, Kong Y, Alexander AC, Albertyn ZI, Boycott KM, Bulman DE, Gordon PM, Innes AM, Knoppers BM, Majewski J, Marshall CR, Parboosingh JS, Sawyer SL, Samuels ME, Schwartzentruber J, Kohane IS, Margulies DM. An international effort towards developing standards for best practices in analysis, interpretation and reporting of clinical genome sequencing results in the CLARITY Challenge. *Genome Biol.*15(3):R53.

10. Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR, Consortium GP. A global reference for human genetic variation. *Nature*. 2015;526(7571):68-74.
11. Naslavsky MS, Yamamoto GL, de Almeida TF, Ezquina SAM, Sunaga DY, Pho N, Bozoklian D, Sandberg TOM, Brito LA, Lazar M, Bernardo DV, Amaro E, Duarte YAO, Lebrão ML, Passos-Bueno MR, Zatz M. Exomic variants of an elderly cohort of Brazilians in the ABraOM database. *Hum Mutat*. 2017;38(7):751-763.
12. Bleichert F, Baserga SJ. The long unwinding road of RNA helicases. *Mol Cell*. 2007;27(3):339-352.
13. Sathya P, Tomkins DJ, Freeman V, Paes B, Nowaczyk MJ. De novo deletion 12q: report of a patient with 12q24.31q24.33 deletion. *Am J Med Genet*. 1999;84(2):116-119.
14. Al-Zahrani J, Al-Dosari N, Abudheim N, Alshidi TA, Colak D, Al-Habit O, Al-Odaib A, Sakati N, Meyer B, Ozand PT, Kaya N. Chromosome 12q24.31-q24.33 deletion causes multiple dysmorphic features and developmental delay: First mosaic patient and overview of the phenotype related to 12q24qter defects. *Mol Cytogenet*. 2011;4:9.
15. Park JP, Graham JM, Andrews PA, Wurster-Hill DH. Ring chromosome 12. *Am J Med Genet*. 1988;29(2):437-440.
16. Jankowsky E, Gross CH, Shuman S, Pyle AM. Active disruption of an RNA-protein interaction by a DExH/D RNA helicase. *Science*. 2001;291(5501):121-125.
17. Lessel D, Schob C, Küry S, Reijnders MRF, Harel T, Eldomery MK, Coban-Akdemir Z, Denecke J, Edvardson S, Colin E, Stegmann APA, Gerkes EH, Tessarech M, Bonneau D, Barth M, Besnard T, Cogné B, Revah-Politi A, Strom TM, Rosenfeld JA, Yang Y, Posey JE, Immken L, Oundjian N, Helbig KL, Meeks N, Zegar K, Morton J, The Ddd Study, Schieving JH, Claasen A, Huentelman M, Narayanan V, Ramsey K, Brunner HG, Elpeleg O, Mercier S, Béziau S, Kubisch C, Kleefstra T, Kindler S, Lupski JR, Kreienkamp HJ, Group CRR. De Novo

- Missense Mutations in DHX30 Impair Global Translation and Cause a Neurodevelopmental Disorder. *Am J Hum Genet.* 2018;102(1):196.
18. Choudhury P, Hackert P, Memet I, Sloan KE, Bohnsack MT. The human RNA helicase DHX37 is required for release of the U3 snoRNP from pre-ribosomal particles. *RNA Biol.* 2019;16(1):54-68.
 19. Mills EW, Green R. Ribosomopathies: There's strength in numbers. *Science.* 2017;358(6363).
 20. Hirata H, Ogino K, Yamada K, Leacock S, Harvey RJ. Defective escape behavior in DEAH-box RNA helicase mutants improved by restoring glycine receptor expression. *J Neurosci.* 2013;33(37):14638-14644.
 21. Tian C, Tan S, Bao L, Zeng Q, Liu S, Yang Y, Zhong X, Liu Z. DExD/H-box RNA helicase genes are differentially expressed between males and females during the critical period of male sex differentiation in channel catfish. *Comp Biochem Physiol Part D Genomics Proteomics.* 2017;22:109-119.
 22. Koopman P, Münsterberg A, Capel B, Vivian N, Lovell-Badge R. Expression of a candidate sex-determining gene during mouse testis differentiation. *Nature.* 1990;348(6300):450-452.
 23. Achermann JC, Domenice S, Bachega TA, Nishi MY, Mendonca BB. Disorders of sex development: effect of molecular diagnostics. *Nat Rev Endocrinol.* 2015;11(8):478-488.
 24. Domenice S, Zamboni Machado A, Moraes Ferreira F, Ferraz-de-Souza B, Marcondes Lerario A, Lin L, Yumie Nishi M, Lisboa Gomes N, Evelin da Silva T, Barbosa Silva R, Vieira Correa R, Ribeiro Montenegro L, Narciso A, Maria Frade Costa E, C Achermann J, Bilharinho Mendonca B. Wide spectrum of NR5A1-related phenotypes in 46,XY and 46,XX individuals. *Birth Defects Res C Embryo Today.* 2016;108(4):309-320.
 25. Suntharalingham JP, Buonocore F, Duncan AJ, Achermann JC. DAX-1 (NR0B1) and steroidogenic factor-1 (SF-1, NR5A1) in human disease. *Best Pract Res Clin Endocrinol Metab.* 2015;29(4):607-619.
 26. Pearlman A, Loke J, Le Caignec C, White S, Chin L, Friedman A, Warr N, Willan J, Brauer D, Farmer C, Brooks E, Oddoux C, Riley B, Shajahan S, Camerino G, Homfray T, Crosby AH, Couper J, David A, Greenfield A, Sinclair A, Ostrer H. Mutations in MAP3K1 cause 46,XY disorders of sex development and implicate a common signal transduction pathway in human testis determination. *Am J Hum Genet.* 2010;87(6):898-904.
 27. Gomes NL, Lerário AM, Machado AZ, Moraes DR, Silva TED, Arnhold IJP, Batista RL, Faria Júnior JAD, Costa EF, Nishi MY, Inacio M, Domenice S, Mendonca BB. Long-term outcomes and molecular analysis of a large cohort of patients with 46,XY disorder of sex development due to partial gonadal dysgenesis. *Clin Endocrinol (Oxf).* 2018.
 28. Sarafoglou K, Ostrer H. Clinical review 111: familial sex reversal: a review. *J Clin Endocrinol Metab.* 2000;85(2):483-493.
 29. Assumpção JG, Benedetti CE, Maciel-Guerra AT, Guerra G, Baptista MT, Scolfaro MR, de Mello MP. Novel mutations affecting SRY DNA-binding activity: the HMG box N65H associated with 46,XY pure gonadal dysgenesis and the familial non-HMG box R30I associated with variable phenotypes. *J Mol Med (Berl).* 2002;80(12):782-790.
 30. Philibert P, Polak M, Colmenares A, Lortat-Jacob S, Audran F, Poulat F, Sultan C. Predominant Sertoli cell deficiency in a 46,XY disorders of sex development

- patient with a new NR5A1/SF-1 mutation transmitted by his unaffected father. *Fertil Steril*. 2011;95(5):1788.e1785-1789.
31. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, Olsson I, Edlund K, Lundberg E, Navani S, Szigartyo CA, Odeberg J, Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist PH, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P, Schwenk JM, Hamsten M, von Feilitzen K, Forsberg M, Persson L, Johansson F, Zwahlen M, von Heijne G, Nielsen J, Pontén F. Proteomics. Tissue-based map of the human proteome. *Science*. 2015;347(6220):1260419.
 32. Thul PJ, Åkesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, Alm T, Asplund A, Björk L, Breckels LM, Bäckström A, Danielsson F, Fagerberg L, Fall J, Gatto L, Gnann C, Hober S, Hjelmare M, Johansson F, Lee S, Lindskog C, Mulder J, Mulvey CM, Nilsson P, Oksvold P, Rockberg J, Schutten R, Schwenk JM, Sivertsson Å, Sjöstedt E, Skogs M, Stadler C, Sullivan DP, Tegel H, Winsnes C, Zhang C, Zwahlen M, Mardinoglu A, Pontén F, von Feilitzen K, Lilley KS, Uhlén M, Lundberg E. A subcellular map of the human proteome. *Science*. 2017;356(6340).
 33. Uhlen M, Zhang C, Lee S, Sjöstedt E, Fagerberg L, Bidkhori G, Benfeitas R, Arif M, Liu Z, Edfors F, Sanli K, von Feilitzen K, Oksvold P, Lundberg E, Hober S, Nilsson P, Mattsson J, Schwenk JM, Brunnström H, Glimelius B, Sjöblom T, Edqvist PH, Djureinovic D, Micke P, Lindskog C, Mardinoglu A, Ponten F. A pathology atlas of the human cancer transcriptome. *Science*. 2017;357(6352).
 34. Jégou B. The Sertoli-germ cell communication network in mammals. *Int Rev Cytol*. 1993;147:25-96.
 35. Skinner MK. Cell-cell interactions in the testis. *Endocr Rev*. 1991;12(1):45-77.
 36. Cheng CY, Mruk DD. Cell junction dynamics in the testis: Sertoli-germ cell interactions and male contraceptive development. *Physiol Rev*. 2002;82(4):825-874.
 37. Rios-Rojas C, Bowles J, Koopman P. On the role of germ cells in mammalian gonad development: quiet passengers or back-seat drivers? *Reproduction*. 2015;149(4):R181-191.
 38. Svingen T, Koopman P. Building the mammalian testis: origins, differentiation, and assembly of the component cell populations. *Genes Dev*. 2013;27(22):2409-2426.

Table 1. Phenotype of 46,XY DSD patients with familial embryonic testicular regression syndrome with rare and predicted pathogenic or likely pathogenic *DHX37*

variants

Nationality		Brazilian		Brazilian		Chilean		Argentinian			Brazilian	
Variables		F1:III-2	F1:III-3	F2:II-4	F2:III-1	F3:II-1	F3:II-2	F4:III-1	F4:III-2	F4:II-4	F5:II-6	F5:III-1
Patient												
Sex of rearing		Male	Male	Male	Female	Male	Male	Male	Male	Male	Female	Male
Age at presentation (yrs)		2.2	1.8	14.0	1.8	0.6	24 days	Newborn	4	2	18	10
Diagnosis		ETRS	ETRS	ETRS	ETRS	ETRS	ETRS	ETRS	ETRS	PGD	PGD	ETRS
External genitalia		Micropenis	Micropenis	Micropenis	Micropenis	Micropenis	Micropenis	Micropenis	Micropenis	Micropenis	Atypical	Micropenis
Gonads		Non-palpable	Non-palpable	Non-palpable	Non-palpable	Non-palpable	Non-palpable	Non-palpable	Non-palpable	Left testis - scrotum	Previous gonadectomy	Non-palpable
Histologic analysis		Small bilateral dysgenetic gonads	Left gonad not found. Right dysgenetic gonad	No gonadal tissue found	Left gonad not found. Small right dysgenetic gonad.	Small bilateral dysgenetic gonads	Small bilateral dysgenetic gonads	No gonadal Tissue	No gonadal tissue	Right gonad not found. Left dysgenetic testis with GCNIS*	Bilateral dysgenetic gonads/	No gonadal tissue
Wolffian derivatives		Present	Present	Present	Present	Present	Present	Present	Present	Present	Present	Present
Mullerian Derivatives	Tubes	Present [¥]	Absent	Present [¥]	Present [¥]	Absent	Present	Present [¥]	Present [¥]	Absent	Absent	Absent
	Uterus	Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent
LH (IU/L)		14.5	12	3.5	1.9	<0.5	<0.5	<0.5	<0.5	26	NA	19
FSH (IU/L)		117	133	87	56	10.9	9.5	NA	9	112	NA	43
Basal Testosterone (ng/dL)		<10	<10	<10	NA	16	<10	38	27	16	NA	21
Testosterone after hCG test (ng/dL)		<10	NA	29	26	18	<10	40	29	NA	NA	NA
Allelic variant		p.Arg308Gln	p.Arg308Gln	p.Arg308Gln	p.Arg308Gln	p.Arg674Trp	p.Arg674Trp	p.Arg674Trp	p.Arg674Trp	p.Arg674Trp	p.Ser595Phe	p.Ser595Phe
Variant state		Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous

NA- not available; PGD- partial gonadal dysgenesis; GCNIS - germ cell neoplasia in-situ; * testicular biopsy; ¥ - Rudimentary Fallopian tubes;

Conversion factors to SI units: T, ng/dL to nmol/L, multiply by 0.0347.

559

560 **Table 2.** Phenotype of 46,XY DSD patients with sporadic gonadal dysgenesis spectrum and heterozygous rare pathogenic or likely pathogenic *DHX37* variants

Nationality		Chinese-American					
Variables		Brazilian					
Patient		F6:II-1	F7:II-1	F8:II-1	F9:II-5	F10:II-1	F11:II-1
Social sex		Male	Male to Female	Female	Female	Male to female	Female
Age at presentation (yrs)		0.18	30	7.7	35	19	3.7
Diagnosis		ETRS	ETRS	PGD	Previous gonadectomy	ETRS	PGD
External genitalia		Micropenis	Micropenis	Female	Previous genitoplasty	Micropenis	Atypical
Gonads		Non-palpable	Non-palpable	Non-palpable	NA	Non-palpable	Non-palpable
Histological analysis		No gonadal tissue	No gonadal tissue	Bilateral dysgenetic gonads	NA	No gonadal tissue	Bilateral dysgenetic gonads
Wolff derivatives		Present	Present	Present	NA	NA	Present
Mullerian derivatives	Tubes	Absent	Present	Absent	NA	Absent	Present
	Uterus	Absent	Absent	Absent	Absent	Absent	Absent
LH (IU/L)		0.1	10	0.1	NA	23	NA
FSH (IU/L)		0.4	40	4.9	NA	62	NA
Basal Testosterone (ng/dL)		<10	NA	<10	NA	21	25
Testosterone after hCG test (ng/dL)		<10	<10	NA	NA	NA	33
Allelic variant		p.Arg308Gln	p.Arg308Gln	p.Arg308Gln	p.Thr304Met	p.Arg674Trp	p.Arg674Trp
Variant state		Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous	Heterozygous

561 NA: not available; PGD- Partial gonadal dysgenesis; Conversion factors to SI units: T, ng/dL to nmol/L, multiply by 0.0347.

562

563 **Table 3.** *In silico* prediction analysis of *DHX37* allelic variants identified in 46,XY DSD patients

Families	Nucleotide changed	AA changed	Functional domain	<i>In silico</i> prediction tools						
				Mutation Taster	Mutation Assessor	SIFT	Polyphen-2	PROVEAN	CADD	GERP
F1, F2, F6, F7, F8	c.923G>A	p.Arg308Gln	Helicase ATP-binding	Disease Cause (score: 0.999)	High functional impact (score: 4.38)	Protein function affected (score 0.001)	Probably damaging (score 1.000)	Deleterious (score -3.93)	35	5.3
F3,F4,F10, F11	c.2020C>T	p.Arg674Trp	Helicase superfamily C-terminal domain	Disease Cause (score 1.000)	Middle functional impact (score: 4.83)	Protein function affected (score 0.001)	Probably damaging (score 1.000)	Deleterious (score -7.42)	33	4.2
F5	c.1784C>T	p.Ser595Phe	Helicase superfamily C-terminal domain	Disease Cause (score: 1.000)	High functional impact (score: 4.26)	Protein function affected (score 0.001)	Benign (score 0.24)	Deleterious (score -5.57)	24.4	4.13
F9	c.911C>T	p.Thr304Met	Helicase ATP-binding	Disease Cause (score 1.000)	High functional impact (score: 4.45)	Protein function affected (score 0.001)	Probably damaging (score 1.000)	Deleterious (score -5.89)	29.8	5.3

564

565 **Table 4.** *DHX37* missense allelic variants identified in 46,XY DSD patients and their frequency in population databases

Families	cDNA position	AA change	Phylogenetic Conservation	State	dbSNP	MAFs in population databases				
						1000 Genomes	ExAC	gnomAD	ABraOm	ESP6500
F1, F2, F6, F7, F8	c.923 G>A	p.Arg308Gln	Highly conserved	Heterozygous	Not available	Absent	Absent	0.00006677 Non-Finnish European	Absent	Absent
F3, F4, F10, F11	c.2020C>T	p.Arg674Trp	Highly conserved	Heterozygous	Not available	Absent	Absent	Absent	Absent	Absent
F5	c.1784C>T	p.Ser595Phe	Highly conserved	Heterozygous	Not available	Absent South Asian	Absent	Absent	Absent	Absent
F9	c.911C>T	p.Thr304Met	Highly conserved	Heterozygous	Not available	Absent	Absent	Absent	Absent	Absent

566
567


568 **Table 5.** Pathogenicity classification of DHX37 variants according to the American College of Medical Genetics and Genomics guidelines

Families	Nucleotide changed	AA changed	Population Data	Computational and prediction data	De novo data	Other data	Classification
F1, F2, F6, F7, F8	c.923G>A	p.Arg308Gln	PM2 ^a	PP2 ^b PP3 ^c	PS2 ^d	PM1 ^e	Pathogenic
F3, F4, F10,F11	c.2020C>T	p.Arg674Trp	PM2 ^a	PP2 ^b PP3 ^c		PM1 ^e	Likely Pathogenic
F5	c.1784C>T	p.Ser595Phe	PM2 ^a	PP2 ^b PP3 ^c		PM1 ^e	Likely Pathogenic
F9	c.911C>T	p.Thr304Met	PM2 ^a	PP2 ^b PP3 ^c		PM1 ^e	Likely Pathogenic

PM2: moderate piece of evidence for pathogenicity; PP3: supporting evidence for pathogenicity by computational (*in silico*) data;
PS2: strong support for pathogenicity when the variants are *de novo*; PP4: supporting evidence using phenotype; PM1: pathogenic moderate;
VUS: Variant of Uncertain Significance.
a Absent from controls (or at extremely low frequency if recessive) in Exome Sequencing Project, 1000 Genomes Project, or Exome Aggregation Consortium.
b Missense variant in a gene that has a low rate of benign missense variation and in which missense variants are a common mechanism of disease
c Multiple lines of computational evidence support a deleterious effect on the gene or gene product (conservation, evolutionary, splicing impact, etc.)
d De novo (both maternity and paternity confirmed) in a patient with the disease and no family history.
e Located in a mutational hot spot and/or critical and well-established functional domain without benign variation.

569
570

Familial cases



Sporadic cases






Figure 1. Pedigrees of the eleven families with potential disease-causing *DHX37* variants. Filled symbols represent affected individuals. The affected males (46,XY males) are indicated by filled squares and the affected individuals raised as females (46,XY females) are shown by large dark dots within the squares. Symbols with a diagonal line represent deceased individuals. The *DHX37* genotype is shown for the individuals whose DNA sample was available; +/- indicates a heterozygous state and -/- indicates a homozygous state for wild-type allele. NA- DNA not available. Paternity and maternity was confirmed in families 6 and 8.

576



577

578 **Figure 2.** The identified variants are localized within conserved helicase domains of *DHX37*. Top: Schematic protein structure of *DHX37* showing conserved
579 motifs of the helicase core region, the helicase associated domain (HA2) and the oligonucleotide/oligosaccharide-binding-fold. Middle: Nucleotide-interacting
580 motifs (I, II, and VI), nucleic acid-binding motifs (Ia, Ib, and IV), motif V, which binds nucleic acid and interacts with nucleotides, and motif III, which couples
581 ATP hydrolysis to RNA unwinding (N- N terminus; C- C terminus). Bottom: Amino acids within conserved motifs of the helicase core region. The position of
582 the first and last amino acid within each motif is denoted below left and right, respectively. The position of the allelic variants identified in this study are
583 indicated with vertical arrows and shown in bold in the different species sequence.



585

586 **Figure 3.** Immunoexpression patterns of DHX37 in testis tissues. **A-** Newborn testis showing
 587 strongly positive staining in occasional spermatogonia (arrow) among numerous Sertoli cells,
 588 some of which show weak cytoplasmic staining (original magnification 100X). **B-**
 589 Seminiferous tubules of a 13 year old boy demonstrating tubules with predominance of Sertoli
 590 cells, all of them negative for DHX37. Note some positive stromal cells (arrow) (original
 591 magnification 100X). **C-** Adult testis of a 54 year old man showing positive Leydig (arrow)
 592 and germ cell staining. **D-** Detail of (C) showing the different pattern of stain in different stages
 593 of germ cells. Note strong perinuclear halo in spermatogonia, progressive paranuclear
 594 condensation in spermatocytes and spermatids, and absence of DHX37 expression in
 595 spermatozoa (original magnification $\times 20X$).

